



Novel Single-Stranded DNA Virus Genomes Recovered from Chimpanzee Feces Sampled from the Mambilla Plateau in Nigeria

Matthew Walters,^a Musa Bawuro,^b Alfred Christopher,^b Alexander Knight,^a Simona Kraberger,^{a,c} Daisy Stainton,^{a,d} Hazel Chapman,^a Arvind Varsani^{a,e,f}

School of the Biological Sciences, University of Canterbury, Christchurch, New Zealand^a; Nigerian Montane Forest Project, Mambilla Plateau, Taraba State, Nigeria^b; Department of Microbiology, Immunology and Pathology, Colorado State University, Fort Collins, Colorado, USA^c; School of Biological Sciences, the University of Queensland, Brisbane, Queensland, Australia^d; Department of Clinical Laboratory Sciences, University of Cape Town, Cape Town, South Africa^e; Biondesign Center for Fundamental and Applied Microbiomics, Center for Evolution and Medicine, School of Life Sciences, Arizona State University, Tempe, Arizona, USA^f

ABSTRACT Metagenomic approaches are rapidly expanding our knowledge of the diversity of viruses. In the fecal matter of Nigerian chimpanzees we recovered three gokushovirus genomes, one circular replication-associated protein encoding single-stranded DNA virus (CRESS), and a CRESS DNA molecule.

Through metagenomic approaches, a plethora of novel viruses have been identified. In the past five years, there has been a significant increase in the identification of single-stranded DNA (ssDNA) viruses in a wide variety of ecosystems and organisms. Animal feces have proven to harbor a rich population of viruses (1–6), and thus we sampled the feces of wild chimpanzees from a small population on the Mambilla Plateau of Nigeria to identify novel ssDNA viruses associated with chimpanzees.

The fecal matter was processed in the same manner as described previously (1–6) and used to generate libraries for Illumina sequencing on an Illumina 2000 platform at Macrogen, Inc. (Republic of Korea). ABYSS version 1.9 was used to *de novo* assemble the paired-end reads. Viral-like sequence contigs that were >750 nucleotides (nt) in length were identified using BLASTx (7). Five contigs were identified with viral-like hits to circular ssDNA viruses. Abutting primers were designed to recover the full viral genomes using KAPA HIFI HotStart DNA polymerase (Kapa Biosystems, USA), and the resulting amplicons were cloned into the pJET2.1 vector (Thermo Fisher Scientific, USA) and Sanger-sequenced by primer-walking at Macrogen, Inc.

Three of the molecules (4,851, 5,016, and 5,184 nt) are most closely related, based on their capsid protein (CP) sequences, to members of the subfamily *Gokushovirinae* in the *Microviridae* family, and we refer to these as Chimpanzee feces-associated microphages (CfaMPs) 1 to 3. Members of *Gokushovirinae* are known to infect obligate intracellular parasites such as *Chlamydia* spp., *Bdellovibrio* spp., and *Spiroplasma* spp. (8). Diverse microvirus-like sequences have been detected in a variety of samples, including marine environments, soil, lakes, peatlands, dragonflies, humans, turkey, methane seep sediments, and bats (9–19). The genome organizations of CfaMP-1 to CfaMP-3 are similar to those of other members of the *Gokushovirinae* family. CfaMP-1 and CfaMP-3 are most closely related, sharing ~82% genome-wide pairwise nucleotide identity, whereas CfaMP-2 shares ~60 to 61% identity with CfaMP-1 and CfaMP-3. The major CP amino acid sequences of CfaMP-1 and CfaMP-3 share 81% identity with each other and 56% identity with the CP-like sequence in the bacterium *Chlamydia trachomatis* (CRH84958) and the Eel River Basin pequenoviruses (KP087936 to KP087957) from cold methane seep sediment (18). The major CP amino acid sequences of CfaMP-2 share

Received 19 December 2016 Accepted 23 December 2016 Published 2 March 2017

Citation Walters M, Bawuro M, Christopher A, Knight A, Kraberger S, Stainton D, Chapman H, Varsani A. 2017. Novel single-stranded DNA virus genomes recovered from chimpanzee feces sampled from the Mambilla Plateau in Nigeria. *Genome Announc* 5:e01715-16. <https://doi.org/10.1128/genomeA.01715-16>.

Copyright © 2017 Walters et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Arvind Varsani, arvind.varsani@asu.edu.

69% identity with isolates from an uncultured bacterium (CDL65862) recovered from a *Rattus norvegicus* isolate (20) and 47% with that of Eel River Basin pequenoviruses.

A circular replication-associated protein (Rep) encoding single-stranded (CRESS) DNA virus (3,811 nt) was also identified, which we refer to as Chimpanzee feces-associated circular DNA virus 1 (CfaV-1). The Rep of CfaV-1 is most closely related (68%) to that of rodent stool-associated circular genome virus (RodSV; JF755404) (21), whereas its putative CP shares 30% identity (58% coverage) with sequences of *Entamoeba histolytica* (EMD46443). A 2,356-nt CRESS DNA molecule was also identified that has a Rep that shares 55% and 51% amino acid identities to those of CfaV-1 and RodSV, respectively.

Diverse ssDNA viruses have been identified in a number of environmental samples, with those identified here further expanding our databases and knowledge of these ssDNA viruses.

Accession number(s). The complete genome sequences of the CfaMP-1, CfaMP-2, and CfaMP-3 microphages have been deposited in GenBank under the accession numbers [KR704913](#), [KR704914](#), and [KR704915](#), respectively. The genome sequences of CfaV-1 and the CRESS DNA molecule were deposited in GenBank under the accession numbers [KR704912](#) and [KR704911](#), respectively.

ACKNOWLEDGMENTS

We thank the Nigerian Montane Forest Project for logistical help and the Taraba State Government and Nigeria National Parks for permission to collect chimpanzee feces.

REFERENCES

- Male MF, Kraberger S, Stainton D, Kami V, Varsani A. 2016. Cycloviruses, gemcircularviruses and other novel replication-associated protein encoding circular viruses in Pacific flying fox (*Pteropus tonganus*) faeces. *Infect Genet Evol* 39:279–292. <https://doi.org/10.1016/j.meegid.2016.02.009>.
- Sikorski A, Argüello-Astorga GR, Dayaram A, Dobson RC, Varsani A. 2013. Discovery of a novel circular single-stranded DNA virus from porcine faeces. *Arch Virol* 158:283–289. <https://doi.org/10.1007/s00705-012-1470-0>.
- Sikorski A, Massaro M, Kraberger S, Young LM, Smalley D, Martin DP, Varsani A. 2013. Novel myco-like DNA viruses discovered in the faecal matter of various animals. *Virus Res* 177:209–216. <https://doi.org/10.1016/j.virusres.2013.08.008>.
- Steel O, Kraberger S, Sikorski A, Young LM, Catchpole RJ, Stevens AJ, Ladley JJ, Coray DS, Stainton D, Dayaram A, Julian L, van Bysterveldt K, Varsani A. 2016. Circular replication-associated protein encoding DNA viruses identified in the faecal matter of various animals in New Zealand. *Infect Genet Evol* 43:151–164. <https://doi.org/10.1016/j.meegid.2016.05.008>.
- Varsani A, Kraberger S, Jennings S, Porzig EL, Julian L, Massaro M, Pollard A, Ballard G, Ainley DG. 2014. A novel papillomavirus in Adelie penguin (*Pygoscelis adeliae*) faeces sampled at the Cape Crozier colony, Antarctica. *J Gen Virol* 95:1352–1365. <https://doi.org/10.1099/vir.0.064436-0>.
- Varsani A, Porzig EL, Jennings S, Kraberger S, Farkas K, Julian L, Massaro M, Ballard G, Ainley DG. 2015. Identification of an avian polyomavirus associated with Adelie penguins (*Pygoscelis adeliae*). *J Gen Virol* 96: 851–857. <https://doi.org/10.1099/vir.0.000038>.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J Mol Biol* 215:403–410. [https://doi.org/10.1016/S0022-2836\(05\)80360-2](https://doi.org/10.1016/S0022-2836(05)80360-2).
- Brentlinger KL, Hafenstein S, Novak CR, Fane BA, Borgon R, McKenna R, Agbandje-McKenna M. 2002. *Microviridae*, a family divided: isolation, characterization, and genome sequence of ϕ MH2K, a bacteriophage of the obligate intracellular parasitic bacterium *Bdellovibrio bacteriovorus*. *J Bacteriol* 184: 1089–1094. <https://doi.org/10.1128/jb.184.4.1089-1094.2002>.
- Labonté JM, Suttle CA. 2013. Previously unknown and highly divergent ssDNA viruses populate the oceans. *ISME J* 7:2169–2177. <https://doi.org/10.1038/ismej.2013.110>.
- Tucker KP, Parsons R, Symonds EM, Breitbart M. 2011. Diversity and distribution of single-stranded DNA phages in the North Atlantic Ocean. *ISME J* 5:822–830. <https://doi.org/10.1038/ismej.2010.188>.
- Reavy B, Swanson MM, Cock PJ, Dawson L, Freitag TE, Singh BK, Torrance L, Mushegian AR, Taliany M. 2015. Distinct circular single-stranded DNA viruses exist in different soil types. *Appl Environ Microbiol* 81: 3934–3945. <https://doi.org/10.1128/AEM.03878-14>.
- Zhong X, Guidoni B, Jacas L, Jacquet S. 2015. Structure and diversity of ssDNA *Microviridae* viruses in two peri-alpine lakes (Annecy and Bourget, France). *Res Microbiol* 166:644–654. <https://doi.org/10.1016/j.resmic.2015.07.003>.
- Roux S, Enault F, Robin A, Ravet V, Personnic S, Theil S, Colombet J, Sime-Ngando T, Debroas D. 2012. Assessing the diversity and specificity of ssDNA *Microviridae* viruses in two peri-alpine lakes (Annecy and Bourget, France). *Res Microbiol* 166:644–654. <https://doi.org/10.1016/j.resmic.2015.07.003>.
- Quaiser A, Dufresne A, Ballaud F, Roux S, Zivanovic Y, Colombet J, Sime-Ngando T, Francez AJ. 2015. Diversity and comparative genomics of *Microviridae* in *Sphagnum*-dominated peatlands. *Front Microbiol* 6:375. <https://doi.org/10.3389/fmicb.2015.00375>.
- Rosario K, Dayaram A, Marinov M, Ware J, Kraberger S, Stainton D, Breitbart M, Varsani A. 2012. Diverse circular ssDNA viruses discovered in dragonflies (Odonata: Epiprocta). *J Gen Virol* 93:2668–2681. <https://doi.org/10.1099/vir.0.045948-0>.
- Minot S, Sinha R, Chen J, Li H, Keilbaugh SA, Wu GD, Lewis JD, Bushman FD. 2011. The human gut virome: inter-individual variation and dynamic response to diet. *Genome Res* 21:1616–1625. <https://doi.org/10.1101/gr.122705.111>.
- Zsak L, Day JM, Oakley BB, Seal BS. 2011. The complete genome sequence and genetic analysis of wca82 a novel uncultured microphage from the turkey gastrointestinal system. *Virol J* 8:331. <https://doi.org/10.1186/1743-422X-8-331>.
- Bryson SJ, Thurber AR, Correa AM, Orphan VJ, Vega Thurber R. 2015. A novel sister clade to the enterobacteria microviruses (family *Microviridae*) identified in methane seep sediments. *Environ Microbiol* 17: 3708–3721. <https://doi.org/10.1111/1462-2920.12758>.
- Li L, Victoria JG, Wang C, Jones M, Fellers GM, Kunz TH, Delwart E. 2010. Bat guano virome: predominance of dietary viruses from insects and plants plus novel mammalian viruses. *J Virol* 84:6955–6965. <https://doi.org/10.1128/JVI.00501-10>.
- Jørgensen TS, Xu Z, Hansen MA, Sørensen SJ, Hansen LH. 2014. Hundreds of circular novel plasmids and DNA elements identified in a rat cecum metagenome. *PLoS One* 9:e87924. <https://doi.org/10.1371/journal.pone.0087924>.
- Phan TG, Kapusinszky B, Wang C, Rose RK, Lipton HL, Delwart EL. 2011. The fecal viral flora of wild rodents. *PLoS Pathog* 7:e1002218. <https://doi.org/10.1371/journal.ppat.1002218>.